## **Supporting information for:**

## The ozone-climate penalty: Past, present, and future

D.J. Rasmussen,<sup>†</sup> Jianlin Hu,<sup>†</sup> Abdullah Mahmud,<sup>‡</sup> and Michael J. Kleeman\*,<sup>†</sup>

Department of Civil and Environmental Engineering, University of California Davis, Davis

CA 95616, and Department of Civil and Environmental Engineering, Portland State

University, Portland OR 97207

E-mail: mjkleeman@ucdavis.edu

Phone: +1 (530) 752-8386. Fax: +1 (530) 752-7872

<sup>\*</sup>To whom correspondence should be addressed

<sup>&</sup>lt;sup>†</sup>Department of Civil and Environmental Engineering, University of California Davis, Davis CA 95616

<sup>&</sup>lt;sup>‡</sup>Department of Civil and Environmental Engineering, Portland State University, Portland OR 97207





Figure S1: Map showing the location of the receptors in the San Joaquin Valley (red circles) and the South Coast Air Basin (purple crosses) where modeled ozone data was extracted from to generate Fig. 4

## A. Ozone-climate penalty (ppb per K); September 8-9, 1993 (10:00-18:00 LDT)






Figure S2: Maps showing the value of the  $O_3$ -climate penalty (ppb  $K^{-1}$ ) for base case  $NO_x$  and VOC emissions (i.e. scaling factors of 1) and the location of the receptors chosen for the isopleth plots (Figs. 2, 3) for (a) the South Coast Air Basin and (b) the San Joaquin Valley